

# Lecture 3: Static Games with Complete Information (Theory)

Albert Banal-Estanol

January 2006

## In the Previous Lecture...

- Game Theory: set of tools to analyse behaviour in the presence of *strategic interdependence*
  - *Extensive form* representation: what players can and cannot do and consequences (tree structure)
  - A *strategy* for a player specifies what to play in each possible circumstance in which the player might be called to play
  - For each *strategy profile* there is an outcome and therefore payoffs
  - *Normal form* representation: players, strategies and payoffs
  - *Mixed and behavioural* strategies: randomisation over the set of pure strategies and over the set of actions in each information set, resp
  - Both types of randomisation are equivalent
-

## Today's Lecture

- What should we expect players to play?
  - Looking for reasonable concepts in simple predictable games and apply these concepts in other settings
  - In this chapter, concentrate in simultaneous move (or "strategic") games
  - Here, a strategy = an action (use normal form)
  - Solution concepts:
    - Use dominant strategies
    - Don't use dominated strategies
    - Play Nash equilibrium strategies (reasonable? existence?)
  - Assume that rationality (and payoffs) are common knowledge:
    - Players are rational and all know that the others are rational and all know that the others know that everyone is rational,...
-

## Dominant Strategies

- Example: "prisoner's dilemma". Two arrested individuals answer separately whether they committed a crime. If both confess, sentence of 5 years in prison each. If none confesses, 2 year each. If one confesses and the other does not, 1 and 10 years, respec. Representation in normal form:

1\2	<i>DC</i>	<i>C</i>
<i>DC</i>	-2,-2	-10,-1
<i>C</i>	-1,-10	-5,-5

- Situations that can be modelled similarly: working in a joint project, arms race,...
- What will be the outcome? Both confessing! Conflict with Pareto-optimality
- Definition: A strategy  $s_i \in S_i$  is a strictly dominant strategy for player  $i$  in game  $\Gamma_N = [I, \{S_i\}, \{u_i(\cdot)\}]$  if for all  $s'_i \neq s_i$  we have

$$u_i(s_i, s_{-i}) > u_i(s'_i, s_{-i}) \text{ for all } s_{-i} \in S_{-i}$$

## Dominated Strategies

- Problem: dominant strategies rarely exist. Examples:

(a)

1\2	L	R
U	1,-1	-1,-1
M	-1,1	1,-1
D	-2,5	-3,2

(b)

1\2	L	R
U	5,1	4,0
M	6,0	3,1
D	6,4	4,4

- Definition: A strategy  $s_i \in S_i$  is a *strictly* dominated strategy for player  $i$  in game  $\Gamma_N = [I, \{S_i\}, \{u_i(\cdot)\}]$  if there exists  $s'_i \in S_i$  such that

$$u_i(s'_i, s_{-i}) > u_i(s_i, s_{-i}) \text{ for all } s_{-i} \in S_{-i}$$

- ...it is *weakly* dominated if there exists  $s'_i \in S_i$  such that

$$u_i(s'_i, s_{-i}) \geq u_i(s_i, s_{-i}) \text{ for all } s_{-i} \in S_{-i} \text{ with } > \text{ for some } s_{-i} \in S_{-i}$$

- Examples: D in game (a) is strictly dominated and U and M in (b) are weakly dominated. Should we rule out weakly dominated strategies as well?
-

## Iterated Elimination of Dominated Strategies

- Example: Modified prisoner's dilemma

1\2	<i>DC</i>	<i>C</i>
<i>DC</i>	0,-2	-10,-1
<i>C</i>	-1,-10	-5,-5

- No dominated strategy for Player 1 but *DC* is dominated for Player 2.
  - If Player 2's strategy is eliminated, then *DC* is dominated for Player 1.
  - Iterative process assuming common knowledge of payoffs and of rationality
  - Results obtained by eliminating iteratively strictly dominated strategies does not depend on the order of deletion. Deleting *weakly* dominated may. Example: in game (b) in previous slide (U,L,M) leads to (D,R) and (M,R,U) leads to (D,L)
  - Extension to mixed strategies (see properties in MWG)
-

## Nash Equilibrium (Nash 1951)

- Definition: A strategy profile  $(s_1, s_2, \dots, s_I)$  constitutes a Nash equilibrium of the game  $\Gamma_N = [I, \{S_i\}, \{u_i(\cdot)\}]$  if for every  $i = 1, \dots, I$

$$u_i(s_i, s_{-i}) \geq u_i(s'_i, s_{-i}) \text{ for all } s'_i \in S_i$$

1\2	<i>l</i>	<i>m</i>	<i>r</i>
<i>U</i>	5,3	0,4	3,5
<i>M</i>	4,0	5,5	4,0
<i>D</i>	3,5	0,4	5,3

- In this example: (M,m) is (the unique) NE
  - Strategies of the NE cannot be strictly dominated
  - In a NE, players have *correct* beliefs about others' play
-

- May still not be unique. Example: Coordination game

$1 \setminus 2$	$ES$	$GC$
$ES$	100,100	0,0
$GC$	0,0	1000,1000

- $(ES, ES)$  and  $(GC, GC)$  are two pure strategy Nash equilibrium
- Definition: A strategy profile  $(\sigma_1, \sigma_2, \dots, \sigma_I)$  constitutes a Nash equilibrium of the game  $\Gamma_N = [I, \{\Delta(S_i)\}, \{u_i()\}]$  if for every  $i = 1, \dots, I$

$$u_i(\sigma_i, \sigma_{-i}) \geq u_i(\sigma'_i, \sigma_{-i}) \text{ for all } \sigma'_i \in \Delta(S_i)$$

- Alternative definition of NE:  $(\sigma_1, \sigma_2, \dots, \sigma_I)$  is a NE iff strategy  $\sigma_i$  is a best response to  $\sigma_{-i}$  for all  $i$
- Definition: In a game  $\Gamma_N = [I, \{\Delta(S_i)\}, \{u_i()\}]$ , a strategy  $\sigma_i$  is a best response for player  $i$  to her rival's strategy  $\sigma_{-i}$ ,  $\sigma_i \in b(\sigma_{-i})$ , if

$$u_i(\sigma_i, \sigma_{-i}) \geq u_i(\sigma'_i, \sigma_{-i}) \text{ for all } \sigma'_i \in \Delta(S_i)$$


---

## A method to Find Pure and Mixed Strategy NE (Example)

1. Find best-response correspondences. For each  $[(p, 1 - p), (q, 1 - q)]$

$$b_1(q) = \begin{cases} 0 & \text{if } q \in [0, \frac{10}{11}) \\ [0, 1] & \text{if } q = \frac{10}{11} \\ 1 & \text{if } q \in (\frac{10}{11}, 1] \end{cases} \quad \text{and} \quad b_2(p) = \begin{cases} 0 & \text{if } p \in [0, \frac{10}{11}) \\ [0, 1] & \text{if } p = \frac{10}{11} \\ 1 & \text{if } p \in (\frac{10}{11}, 1] \end{cases}$$

2. Assume that  $[(p^*, 1 - p^*), (q^*, 1 - q^*)]$  is a MSNE and look for conditions:

If  $p^* = 0$  then  $q^* = 0$ . If  $q^* = 0$  then  $p^* = 0$ . Hence  $[(0, 1), (0, 1)]$  is a NE

Similarly  $[(1, 0), (1, 0)]$  is a NE.

If  $0 < p^* < 1$  then  $q^* = \frac{10}{11}$  and then since  $0 < q^* < 1$ ,  $p^* = \frac{10}{11}$ .  
 $[(\frac{10}{11}, \frac{1}{11}), (\frac{10}{11}, \frac{1}{11})]$  is a NE

- Notice that (here and in general) in a MSNE, each player is indifferent among all the pure strategies played with positive probability

## Existence

- Proposition: Every game  $\Gamma_N = [I, \{\Delta(S_i)\}, \{u_i()\}]$  in which the sets  $S_i$  are finite, has a mixed strategy NE.
  - Proposition: A NE in a game  $\Gamma_N = [I, \{S_i\}, \{u_i()\}]$  exists if for all  $i = 1, \dots, I$  :
    - a)  $S_i$  is a non-empty, convex, and compact set of some Euclidean space  $\mathbb{R}^M$
    - b)  $u_i(s_1, \dots, s_I)$  is continuous in  $(s_1, \dots, s_I)$  and quasiconcave in  $s_i$
-